arXiv:cond-mat/0612165v3 [cond-mat.str-el] 5 May 2007

Coexistence of vector chiral order and Tomonaga-Luttinger liquid in the frustrated three-leg spin tube in a magnetic field

Masahiro Sato
Synchrotron Radiation Research Unit, Japan Atomic Energy Agency, Sayo, Hyogo 679-5148, Japan
CREST Japan Science and Technology Agency, Kawaguchi, Saitama 332-0012, Japan

(Dated: February 6, 2008)

The frustrated three-leg antiferromagnetic spin-$\frac{1}{2}$ tube with a weak interchain coupling in a magnetic field is investigated by means of Abelian bosonization techniques. It is clearly shown that a vector chiral long-range order and a one-component Tomonaga-Luttinger liquid coexist in a wide magnetic-field region from a state with a small magnetization to a nearly saturated one. The chiral order is predicted to still survive in the intermediate plateau state. We further predict that (even) when the strength of one bond in the three rung couplings is decreased (increased), an Ising-type quantum phase transition takes place and the chirality vanishes (no singular phenomena occur and the chiral order is maintained). Even without magnetic fields, the chiral order would also be present if the spin tube possesses easy-plane anisotropy.

PACS numbers: 75.10.Jm, 75.40.Cx, 75.50.Ee

I. INTRODUCTION

Quantum spin tubes−namely, systems of coupled spin chains with a *periodic boundary condition* (PBC) for the interchain (rung) direction−have been studied since late 1990s.[1,2,3,4,5,6,7,8,9,10,11] In particular, odd-leg spin tubes have attracted much interest, because the antiferromagnetic (AF) rung coupling yields frustration. It has been recognized now that coupled spin chains including spin tubes offer fairly richer physics compared with single spin chains.[12] Recently, a few new spin-tube compounds such as a three-leg tube[6] $[(\mathrm{CuCl_2tachH})_3\mathrm{Cl}]\mathrm{Cl}_2$ and a nine-leg one[5] $\mathrm{Na_2V_3O_7}$ have been fabricated. The presence of these materials has further promoted the study of spin tubes.

For *nonfrustrated* AF spin-S ladders−namely, coupled AF chains with an *open boundary condition* for the rung−the following universal feature (generalization of Haldane's prediction) has been elucidated:[13,14,15] if both $2S$ and the number of chains, N, are odd, the low-energy physics is described by a gapless, one-component Tomonaga-Luttinger liquid (TLL) and no symmetry breakings take place, whereas for all the other cases ($2S\times N=$ even), the system has a gapped spectrum with conserving all symmetries. This even-odd property, however, does not always hold in *frustrated* coupled spin chains. In the three-leg AF spin-$\frac{1}{2}$ tube where the rung triangle induces geometrical frustration, it is known[1] that the one-site translational symmetry along the chain direction is spontaneously broken and a finite gap between a ground state and the lowest excitation exists.

In this paper, we point out another striking difference between the three-leg nonfrustrated ladder and the frustrated tube: namely, utilizing bosonization techniques, we show in a clear way that a vector chiral long-range order resides in the magnetic-field-induced TLL phase of the three-leg AF spin-$\frac{1}{2}$ tube at least in the weak-rung-coupling regime. In the chiral phase, the parity symmetry for the rung direction is spontaneously broken. On the contrary, the chiral order is absent in the corresponding ladder. For one-dimensional gapped $U(1)$-symmetric coupled spin chains, (i) the field-induced magnon-condensed gapless state is usually described by a standard TLL and conserves all symmetries and (ii) many physicists have focused only on whether nontrivial magnetization plateau states exist or not. Therefore, the possibility of chirality in spin tubes has not been considered well so far.[16] We also show that the chiral order continues in the intermediate plateau state of the spin tube. These predictions means that the three-leg spin tube possesses both gapless and gapped chiral-ordered states. We further discuss a rung-coupling modification in the three-leg spin tube: the strength of one bond of three rung couplings is changed. If the strength of the changed bond is extremely increased (decreased), the system approaches a two-leg ladder plus single chain (a three-leg ladder). Both limits possess no geometrical frustration. Interestingly, it is found that in the case of increasing the bond strength, the chiral order always remains unbroken, but an Ising-type quantum phase transition takes place and the chirality disappears at the Ising critical point in the other case.

The organization of the paper is as follows. In Sec. II, we define a spin tube model and briefly explain the bosonization method. The next two sections are the main part of the paper. We show the mechanism of the vector chiral order in Sec. III. Section IV is devoted to investigating effects of the rung deformation. Finally, conclusions and brief discussions about our results are presented in Sec. V.

II. MODEL AND METHOD

In this paper, we mainly consider the three-leg AF spin-$\frac{1}{2}$ tube with a homogeneous rung coupling. The

Hamiltonian is defined as

$$\mathcal{H}=\sum_{n=1}^{3}\sum_{j}\left[J\vec{S}_{n,j}\cdot\vec{S}_{n,j+1}+J_{\perp}\vec{S}_{n,j}\cdot\vec{S}_{n+1,j}-HS^{z}_{n,j}\right],\quad(1)$$

where $\vec{S}_{n,j}$ is spin-$\frac{1}{2}$ operator on site j of the nth chain (n = 1-3), $J > 0$ ($J_\perp > 0$) is the AF intrachain (rung) coupling, and the PBC $\vec{S}_{4,j}=\vec{S}_{1,j}$ is imposed. We begin with the independent three chains in a field H. From Abelian bosonization,[12] the nth chain in the low-energy limit is mapped to a Gaussian model $\mathcal{H}_n=\int dx\ \frac{v}{2}\left[K(\partial_x\theta_n)^2+K^{-1}(\partial_x\phi_n)^2\right]$, where ϕ_n is the scalar boson field and θ_n is the dual to ϕ_n. The TLL parameter K and the spin-wave velocity v depend on J and H; when H is changed from 0 to the saturation field $2J$, v (K) monotonically decreases (increases) from $\pi Ja_0/2$ (1/2) to 0 (1) [a_0: lattice constant]. The spin operators are also bosonized as

$$\begin{aligned}S^{z}_{n,j} &\approx M+a_0\partial_x\phi_n/\sqrt{\pi}\\ &\quad+(-1)^{j}A_1\sin(\sqrt{4\pi}\phi_n+2\pi Mj)+\cdots,\quad(2a)\end{aligned}$$

$$\begin{aligned}S^{+}_{n,j} &\approx e^{i\sqrt{\pi}\theta_n}\big[(-1)^{j}B_0\\ &\quad+B_1\sin(\sqrt{4\pi}\phi_n+2\pi Mj)+\cdots\big],\quad(2b)\end{aligned}$$

where $M(H)=\langle S^{z}_{n,j}\rangle$, and A_l and B_l are nonuniversal constants of $O(1)$.[17,18] This formula indicates that the period of ϕ_n (θ_n) is $\sqrt{\pi}$ ($\sqrt{4\pi}$). Using $\mathcal{H}_n$ and formula (2), we can straightforwardly derive the following effective Hamiltonian of the model (1) in the weak-rung-coupling regime:

$$\begin{aligned}\mathcal{H}_{\rm eff}=\int dx\ &\sum_{n=1}^{3}\frac{v}{2}\left[K(\partial_x\theta_n)^2+K^{-1}(\partial_x\phi_n)^2\right]\\ &+\frac{2MJ_\perp}{\sqrt{\pi}}\sum_n\partial_x\phi_n+\frac{J_\perp a_0}{\pi}\sum_n\partial_x\phi_n\partial_x\phi_{n+1}\\ &+\sum_n J_\perp a_0^{-1}\Big[B_0^2\cos(\sqrt{\pi}(\theta_n-\theta_{n+1}))\\ &+\frac{A_1^2}{2}\cos(\sqrt{4\pi}(\phi_n-\phi_{n+1}))\Big]+\cdots,\quad(3)\end{aligned}$$

where $\phi_4=\phi_1$, $\theta_4=\theta_1$, and we have written only the important part among all the rung-coupling terms and neglected terms with oscillating factors $e^{li\pi Mj}$ or $(-1)^j$, which are irrelevant. In the bosonization picture, symmetries of the spin tube (1) are represented as follows: a $U(1)$ rotation around the S^z axis $S^{+}_{n,j}\to e^{i\gamma}S^{+}_{n,j}$, the one-site translation along the chain $S^{\alpha}_{n,j}\to S^{\alpha}_{n,j+1}$, that along the rung $S^{\alpha}_{n,j}\to S^{\alpha}_{n+1,j}$, the site-parity transformation for the chain $S^{\alpha}_{n,j}\to S^{\alpha}_{n,-j}$, and that for the rung $S^{\alpha}_{1,j}\leftrightarrow S^{\alpha}_{3,j}$, respectively, correspond to $\theta_n\to\theta_n+\gamma/\sqrt{\pi}$, $(\phi_n,\theta_n)\to(\phi_n+\sqrt{\pi}(M+1/2),\theta_n+\sqrt{\pi})$, $(\phi_n,\theta_n)\to(\phi_{n+1},\theta_{n+1})$, $(\phi_n(x),\theta_n(x))\to(-\phi_n(-x)+\sqrt{\pi}/2,\theta_n(-x))$ and $(\phi_1,\theta_1)\leftrightarrow(\phi_3,\theta_3)$. These symmetries strongly restrict possible terms in $\mathcal{H}_{\rm eff}$. In other words, the symmetries reduce the number of coupling constants. As a result, for all vertex operators without oscillating factors and conformal spin, only $\sum_n\cos[l\sqrt{4\pi}(\phi_n-\phi_{n+1})]$ and $\sum_n\cos[l\sqrt{4\pi}(\theta_n-\theta_{n+1})]$ are allowed to exist in Eq. (3). The most relevant terms with $l=1$ indeed appear in Eq. (3). Utilizing the effective theory (3), we study the low-energy properties of the spin tube below. We stress here that almost all the discussions below are applicable to other three-leg spin tubes with symmetry-conserved perturbations (XXZ anisotropy, further-neighbor interactions, etc.).



FIG. 1: Relationship between $\{\theta_n\}$ and $\{\Theta_q\}$. C is a constant.

III. VECTOR CHIRAL ORDER

Let us introduce new boson fields $\Phi_0=\sum_n\phi_n/\sqrt{3}$, $\Phi_1=(\phi_1-\phi_3)/\sqrt{2}$, $\Phi_2=(\phi_1+\phi_3-2\phi_2)/\sqrt{6}$, $\Theta_0=\sum_n\theta_n/\sqrt{3}$, $\Theta_1=(\theta_1-\theta_3)/\sqrt{2}$, and $\Theta_2=(\theta_1+\theta_3-2\theta_2)/\sqrt{6}$. The relationship between old and new bosons is illustrated in Fig. 1. As one will see later, these fields help us to detect a vector chiral order. Using these new fields, we can diagonalize the boson bilinear part in Eq. (3). Consequently, the effective Hamiltonian is rewritten as

$$\begin{aligned}\mathcal{H}_{\rm eff}\ =\ \int dx\ &\sum_{q=0}^{2}\frac{v_q}{2}\left[K_q(\partial_x\Theta_q)^2+K_q^{-1}(\partial_x\Phi_q)^2\right]\\ &+2\sqrt{\frac{3}{\pi}}MJ_\perp\partial_x\Phi_0+B_0^2\frac{J_\perp}{a_0}V[\Theta_1,\Theta_2]\\ &+A_1^2\frac{J_\perp}{2a_0}V[\Phi_1,\Phi_2]+\cdots,\quad(4)\end{aligned}$$

where the potential $V[\alpha,\beta]$ is defined as

$$V=2\cos\left(\sqrt{\frac{\pi}{2}}\alpha\right)\cos\left(\sqrt{\frac{3\pi}{2}}\beta\right)+\cos\left(\sqrt{2\pi}\alpha\right).\quad(5)$$

The new TLL parameters K_q and velocities v_q are evaluated as

$$\begin{aligned}K_0=Kf_0^{-1},\quad & v_0=vf_0,\\ K_{1,2}=Kf_g^{-1}=K_g,\quad & v_{1,2}=vf_g=v_g,\end{aligned}\quad(6)$$

where $f_0=(1+\frac{2K}{\pi}\frac{J_\perp a_0}{v})^{1/2}$ and $f_g=(1-\frac{K}{\pi}\frac{J_\perp a_0}{v})^{1/2}$. One finds that the Hamiltonian (4) does not contain



FIG. 2: (a) Potential $V[\Theta_1, \Theta_2]$. (b) Physically relevant zone (enclosed by the green line) in the projected Θ_1-Θ_2 plane. $\vec{K}_n$ play the role of the primitive translation vectors. The rung-parity transformation causes $\Theta_1 \to -\Theta_1$, and the rung translation by one site does $\vec{K}_n \to \vec{K}_{n+1}$.

any vertex operator with Φ_0 or Θ_0. This is strongly supported by symmetries of the tube. The Φ_0 sector hence provides a one-component TLL. The derivative term $\partial_x \Phi_0$ can be absorbed into the Gaussian part and yields only a small correction of M. On the other hand, the remaining $\Phi_{1,2}$ sectors are subject to the effect of the potential V. Since the scaling dimensions of $V[\Phi_1, \Phi_2]$ and $V[\Theta_1, \Theta_2]$ are, respectively, evaluated as $2K_g$ and $1/(2K_g)$, the latter $V[\Theta_1, \Theta_2]$ is always dominant and relevant [i.e., $1/(2K_g) < 2$] from a small M to the saturation. The phase fields Θ_1 and Θ_2 thus are pinned at the minimum of $V[\Theta_1, \Theta_2]$, and $\Phi_{1,2}$ sectors have massive spectra. Consequently, the low-energy physics of the tube (1) is governed by a TLL in the Φ_0 sector. The TLL just corresponds to the known field-induced critical phase.[3] [For the case of $M = 0$, $e^{li\pi Mj} = 1$ and relevant vertex operators with Φ_0 appear in Eq. (4). They are expected to cause the Φ_0 sector to obtain a gapped spectrum.]

The scenario leading to the TLL here is already known well.[3,19] However, as one will see below, characteristics of the frustrated spin tube (1) are hidden in the form of the potential V. Figure 2(a) and Eq. (5) show that six minimum points $(\Theta_1, \Theta_2) = (\pm\sqrt{2\pi}/3, \pm\sqrt{2\pi/3})$ and $(\pm 2\sqrt{2\pi}/3, 0)$ lie around the origin $(0, 0)$. [Inversely, $(0, 0)$ is the unique minimum of the potential $J_\perp V$ for $J_\perp < 0$.] Not all, however, are physically meaningful, because phase fields θ_n have a period. Projecting the physically relevant cubic space $\theta_n \in [-\sqrt{\pi}, \sqrt{\pi}]$ (mod $\sqrt{4\pi}$) onto the Θ_1-Θ_2 plane (see Fig. 1), we obtain a diamond zone as in Fig. 2(b). Therefore, it is enough to consider only two minimum points $(\Theta_1^{\pm}, \Theta_2) = (\pm\sqrt{2\pi}/3, \sqrt{2\pi/3})$ that satisfy $\Theta_1^+ = -\Theta_1^- \neq 0$. Since the rung-parity transformation $(\phi_1, \theta_1) \leftrightarrow (\phi_3, \theta_3)$ causes $\Theta_1 \to -\Theta_1$, pinning Θ_1 to these two minima implies the spontaneous breakdown of the rung-parity symmetry. As a candidate of the rung-parity order parameter, we can propose a vector chirality $\vec{\kappa}_{n,j} = \vec{S}_{n,j} \times \vec{S}_{n+1,j}$, which changes sign via the rung-parity operation. From the formula (2), the z component of a chirality is evaluated as

$$\langle \kappa_{3,j}^z \rangle \approx -B_0^2 \left\langle \sin\left(\sqrt{2\pi}\Theta_1\right) \right\rangle + \cdots = \mp \,\text{finite}, \qquad (7)$$

for $\Theta_1^{\pm}$. Remarkably, the leading term of the chirality does not contain the massless fields (Φ_0, Θ_0). Similarly, $\langle \kappa_{1(2),j}^z \rangle$ are shown to be equivalent to $\langle \kappa_{3,j}^z \rangle$. We thus conclude that the vector chiral long-range order exists in the field-induced TLL phase. The chirality correlation function exhibits an exponential decay: $\langle \kappa_{3,j}^z \kappa_{3,0}^z \rangle \approx \langle \kappa_{3,j}^z \rangle^2 + C_0 e^{-|j|/\xi}/\sqrt{|j|} + \cdots$. Since any artificial approximation such as a mean-field decoupling is not used to obtain Eq. (7), the prediction of the chiral order is highly reliable. Note that in the chiral phase, the spontaneous breakdown of the Z_2 rung-parity symmetry occurs, but the $U(1)$ spin-rotational symmetry is preserved.

For the spin-$\frac{1}{2}$ tube (1), it is predicted that the intermediate plateau state appears at $M = 1/6$ in the region $J_\perp > J_\perp^c (\sim 0.1J)$.[3] In the bosonization picture, the spin gap of the plateau state is interpreted as that of the Φ_0 sector, which is induced by a relevant higher-order umklapp term $\cos(4\sqrt{3\pi}\Phi_0 + 12\pi Mj) = \cos(4\sqrt{3\pi}\Phi_0)$. On the other hand, the $\Phi_{1,2}$ sectors are hardly influenced by whether the Φ_0 sector obtains a gapped spectrum or not. Therefore, we predict that the chiral order still continues in the intermediate plateau phase at least if $J_\perp (> J_\perp^c)$ is sufficiently small. This chiral plateau state is reminiscent of the narrow chiral phase in the classical AF XY model on a triangular lattice.[20] It is noteworthy that by controlling the strength of the magnetic field, one can realize both *gapless and gapped* chiral states in *single* spin tube.

IV. RUNG DEFORMATION

In this section, we discuss the rung deformation. Let us modify the coupling between the first and third chains as

$$J_\perp \vec{S}_{1.j} \cdot \vec{S}_{3,j} \;\to\; J_\perp (1+\delta) \vec{S}_{1.j} \cdot \vec{S}_{3,j}. \qquad (8)$$

This explicitly violates the rung translational symmetry, but conserves the parity symmetry between the first and third chains. Therefore, $\langle \kappa_{3,j}^z \rangle$ is still valid as a rung-parity order parameter. The cases of $\delta = -1$ and $\delta \to +\infty$, respectively, correspond to a three-leg ladder and a system of a two-leg ladder plus single chain. Note here that if we set $J_\perp \delta = \text{const} \ll J$, the present weak-rung-coupling approach is available even for the case of a large $|\delta| \gg 1$. A finite δ brings new bosonic terms $\delta J_\perp M \partial_x(\phi_1 + \phi_3)/\sqrt{\pi}$, $\delta J_\perp a_0 \partial_x \phi_1 \partial_x \phi_3/\pi$, $\delta J_\perp B_0^2 a_0 \cos(\sqrt{2\pi}\Theta_1)$, etc. The first term can be absorbed into the Gaussian part via the shift of Φ_q, which does not affect dual fields Θ_q and yields a small deviation from the uniform magnetization, $\langle S_{2,j}^z \rangle \neq \langle S_{1(3),j}^z \rangle$. It is not surprising because (as mentioned above) δ breaks the rung translational symmetry. The second boson bilinear term is expected not to qualitatively influence the low-energy physics. Actually, introducing a new basis different from (Φ_q, Θ_q), we can diagonalize all the bilinear



FIG. 3: Deformed potential $V_\delta[\Theta_1, \Theta_2]$.

terms. The main effect of the rung modification originates from the third cosine term which varies the form of the potential $V[\Theta_1, \Theta_2]$ as follows:

$$V[\Theta_1, \Theta_2] \to V_\delta[\Theta_1, \Theta_2] = \tag{9}$$
$$2\cos\left(\sqrt{\frac{\pi}{2}}\Theta_1\right)\cos\left(\sqrt{\frac{3\pi}{2}}\Theta_2\right) + (1+\delta)\cos\left(\sqrt{2\pi}\Theta_1\right).$$

The deformed potential V_δ is illustrated in Fig. 3. From V_δ, one finds that when δ increases [decreases and reaches $\delta_{\rm cl} = -0.5$], the two minimum points still survive with keeping $\Theta_1 \neq 0$ [approach with each other and meet at $(\Theta_1, \Theta_2) = (0, \sqrt{2\pi/3})$]. The chirality $\langle \kappa^z_{3,j} \rangle$ is hence shown to be always finite when an arbitrary positive δ is applied. In other words, once an infinitesimal exchange is added between a two-leg AF ladder and single chain, a vector chiral order immediately emerges. On the other hand, for the case of $\delta < 0$, the vector chiral order is predicted to vanish at a certain point $\delta = \delta_c$ and the low-energy physics is described by the TLL of the Φ_0 sector for $\delta < \delta_c$. The true value of δ_c must be renormalized from the classical value $\delta_{\rm cl}$ by the effect of the Gaussian part and must depend on $J_\perp/J$ and H/J.

In order to further understand the physics near the phase transition at $\delta = \delta_c$, it is necessary to go beyond the semiclassical analysis of V_δ. If $\cos(\sqrt{3\pi/2}\Theta_2)$ in V_δ is allowed to be replaced with its expectation value, the effective Hamiltonian for the Φ_1 sector is written as the following double sine-Gordon (dSG) model[21,22]:

$$\mathcal{H}_{[\Phi_1,\Theta_1]} \approx \int dx\ \left[K_g(\partial_x\Theta_1)^2 + K_g^{-1}(\partial_x\Phi_1)^2\right]$$
$$-\frac{J_\perp}{a_0}B_0^2\left[C_1\cos\left(\sqrt{\frac{\pi}{2}}\Theta_1\right) - (1+\delta)\cos\left(\sqrt{2\pi}\Theta_1\right)\right], \tag{10}$$

where $C_1 = \langle\cos(\sqrt{3\pi/2}\Theta_2)\rangle$. It is believed that for $K_g > 1/4$, the dSG model exhibits a second-order Ising transition by tuning the coupling constants. Indeed, as δ passes through a critical value δ_c, the potential of the dSG model changes from a double-well type to a single-bottom one. We thus conclude that the transition at $\delta = \delta_c$ is in the Ising universality class. The semiclassical analysis of V_δ becomes more reliable with V_δ being more relevant. It is hence expected that the value δ_c monotonically decreases and approaches $\delta_{\rm cl}$ when $K_g(\approx K)$ increases—i.e., when H becomes close to the upper critical field. Equation (3.11) and Fig. 12 in Ref. 22 also support this prediction.



FIG. 4: Phase diagram of the spin-$\frac{1}{2}$ tube. $J_\perp(< J_\perp^c)$ is fixed in the weak-rung-couping regime. The bosonization approach is less valid near both the lower and the upper critical fields. The upper critical field here is evaluated by the energy dispersion of one magnon.

Since $\langle\kappa^z_{3,j}\rangle \propto \langle\sin(\sqrt{2\pi}\Theta_1)\rangle \approx \sqrt{2\pi}\langle\Theta_1\rangle$ holds near the transition, the chirality would play the role of the Ising order parameter. Comparing our dSG model (10) and results [Eqs. (29) and (58)] in Ref. 21, we can find

$$\langle\kappa^z_{3,j}\rangle \sim (\delta - \delta_c)^{1/8}, \tag{11}$$

near $\delta \to \delta_c + 0$. [The mean-field analysis of V_δ leads to $\langle\kappa^z_{3,j}\rangle \sim (\delta - \delta_c)^{1/2}$.] The chirality correlation function behaves as $\langle\kappa^z_{3,j}\kappa^z_{3,0}\rangle \sim C_2/|j|^{1/4}$ just at the critical point. Moreover, near the point, it follows the results in Ref. 23.

Sakai *et al.* have recently predicted that when δ is added to the spin tube in *zero field*, the spin gap rapidly disappears.[24] Combining this prediction and ours, we construct the phase diagram on the δ-H plane as in Fig. 4.

V. CONCLUSIONS AND DISCUSSIONS

We have studied the three-leg AF spin-$\frac{1}{2}$ tube (1) with a weak rung coupling under a magnetic field making use of Abelian bosonization techniques. In Sec. III, we have definitely elucidated without the help of any artificial approximation that the vector chiral long-range order occurs in the magnetic-field-driven one-component TLL phase. Due to the chiral order, the discrete rung-parity symmetry is spontaneously broken, but the $U(1)$ spin-rotational symmetry remains unbroken. We have also been predicted that the chiral order survives even in the intermediate plateau state with $M = 1/6$. It is remarkable that one can obtain both gapless and gapped chiral states by changing the strength of the magnetic field. In Sec. IV, we have investigated effects of the rung deformation in Eq. (8). It has been shown that when a positive δ is introduced with fixing $J_\perp\delta$ the chiral order and the TLL always continue, whereas when δ is decreased the chiral order vanishes at a certain Ising critical point $\delta = \delta_c < 0$ and then a standard TLL phase appears (see Fig. 4). This implies that vector chirality correlations are favored in two-leg spin ladders[25] as well as in three-leg tubes.

Quite recently, based on the spin-wave picture, we have predicted that the same chiral order still survives a certain regime in the vicinity of saturation of the spin tube with $\delta = 0$.[26] Therefore, it is inferred that the tube possesses the chiral order in a quite wide range of the magnetic field. One immediately finds that the classical three-leg spin tube always has a vector chiral order from the zero field to the upper critical one. We thus may say that the classical nature is strong in the weak-rung-coupling area of the quantum spin tube (1). Actually, as well known, the AF spin-$\frac{1}{2}$ chain, which is the starting point of our analysis, has a large instability toward a classical Néel ordering.

The discussion in this paper tells us that even without magnetic fields, a vector chiral order could also occur if a spin tube consists of three spin chains with a large $K(> 1/2)$: for instance, the condition $K > 1/2$ is satisfied in a chain with easy-plane XXZ anisotropy. It is predicted that chiral orders exist in a certain parameter regime of easy-plane zigzag spin chains.[27] Our results also suggests that a field-induced vector chiral order can appear in other frustrated odd-leg spin tubes−i.e., five-, seven-, nine-leg tubes, etc., tubes.

Finally, we briefly mention the possibility of experimentally detecting the vector chirality. The chiral order predicted in this paper must be destroyed by an effect of thermal fluctuation. However, chirality correlations are expected to still be strong if the temperature is sufficiently low. Moreover, weak three-dimensional interactions among tubes can stabilize the long-range chiral order: the gapless chiral order in the tube is expected to change into a conventional umbrella spin structure due to the interactions. Four-point spin correlation functions include features of the above chiral order or strong chiral correlation. Such functions, in principle, are measured in polarized neutron scattering, electromagnetic-wave resonance, etc.

Acknowledgments

This work is supported by a Grant-in-Aid for Scientific Research (B) (No. 17340100) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

---

[1] K. Kawano and M. Takahashi, J. Phys. Soc. Jpn. **66**, 4001 (1997).

[2] H. J. Schulz, cond-mat/9605075.

[3] D. C. Cabra, A. Honecker, and P. Pujol, Phys. Rev. Lett. **79**, 5126 (1997); Phys. Rev. B **58**, 6241 (1998).

[4] R. Citro, E. Orignac, N. Andrei, C. Itoi, and S. Qin, J. Phys.: Condens. Matter **12**, 3041 (2000).

[5] P. Millet, J. Y. Henry, F. Mila and J. Galy, J. Solid State Chem. **147**, 676 (1999).

[6] J. Schnack, H. Nojiri, P. Kögerler, G. J. T. Cooper and L. Cronin, Phys. Rev. B **70**, 174420 (2004).

[7] A. Lüscher, R. M. Noack, G. Misguich, V. N. Kotov and F. Mila, Phys. Rev. B **70**, 060405(R) (2004).

[8] J.-B. Fouet, A. Läuchli, S. Pilgram, R. M. Noack and F. Mila, Phys. Rev. B **73**, 014409 (2006).

[9] K. Okunishi, S. Yoshikawa, T. Sakai, and S. Miyashita, Prog. Theor. Phys. Supp. **159**, 297 (2005).

[10] M. Sato, Phys. Rev. B **72**, 104438 (2005).

[11] M. Sato and M. Oshikawa, Phys. Rev. B **75**, 014404 (2007).

[12] For example see A. O. Gogolin, A. A. Nersesyan and A. M. Tsvelik, *Bosonization and Strongly Correlated Systems* (Cambridge University Press, Cambridge, England, 1998); T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, New York, 2004).

[13] G. Sierra, J. Phys. A **29**, 3299 (1996); Lect. Notes Phys. **478**, eds. G. Sierra and M. A.Martin-Delgado (Springer-Verlag, 1997); see also cond-mat/9610057.

[14] A. G. Rojo, Phys. Rev. B **53**, 9172 (1996).

[15] S. Dell'Aringa, E. Ercolessi, G. Morandi, P. Pieri, and M. Roncaglia, Phys. Rev. Lett. **78**, 2457 (1997).

[16] In this paper, we discuss a vector chiral order with *spontaneously* breaking the rung-parity symmetry in a spin tube. Note, however, that there is another mechanism of a vector chiral order induced by a rung-parity-breaking three-spin chiral term which can emerge via a virtual process on an electron tube system under a magnetic field. Of course, a spin tube Hamiltonian with such chiral terms does not possess the rung-parity symmetry, and the resulting finite expectation value of vector chirality does not mean any spontaneous symmetry breaking. For more detailed discussions, see D. Sen and R. Chitra, Phys. Rev. B **51**, 1922 (1995); R. Chitra and R. Citro, Phys. Rev. B **63**, 054441 (2001).

[17] S. Lukyanov and A. Zamalodchikov, Nucl. Phys. B **493**, 571 (1997).

[18] T. Hikihara and A. Furusaki, Phys. Rev. B **69**, 064427 (2004).

[19] M. Oshikawa, M. Yamanaka, and I. Affleck, Phys. Rev. Lett. **78**, 1984 (1997).

[20] For example see S. Miyashita and H. Shiba, J. Phys. Soc. Jpn. **53**, 1145 (1984); S. Lee and K. C. Lee, Phys. Rev. B **57**, 8472 (1998); L. Capriotti, R. Vaia, A. Cuccoli and V. Tognetti, Phys. Rev. B **58**, 273 (1998).

[21] M. Fabrizio, A. O. Gogolin and A. A. Nersesyan, Nucl. Phys. B **580**, 647 (2000).

[22] Z. Bajnok, L. Palla, G. Takács and F. Wágner, Nucl. Phys. B **601**, 503 (2001).

[23] T. T. Wu, B. M. McCoy, C. A. Tracy, and E. Barouch,

Phys. Rev. B **13**, 316 (1976).

[24] T. Sakai, M. Matsumoto, K. Okunishi, K. Okamoto, and M. Sato, Physica E **29**, 633 (2005).

[25] M. Sato, cond-mat/0703396 (unpublished).

[26] M. Sato and T. Sakai, Phys. Rev. B **75**, 014411 (2007).

[27] A. A. Nersesyan, A. O. Gogolin and F. H. L. Eßler, Phys. Rev. Lett. **81** 910 (1998); M. Kaburagi, H. Kawamura and T. Hikihara, J. Phys. Soc. Jpn. **68**, 3185 (1999); T. Hikihara, M. Kaburagi, and H. Kawamura, Phys. Rev. B **63**, 174430 (2001); A. K. Kolezhuk, Phys. Rev. B **62**, R6057 (2000); P. Lecheminant, T. Jolicoeur, and P. Azaria, Phys. Rev. B **63**, 174426 (2001).